

# Music Therapy for Tinnitus Patients- A Prospective Pilot Study

Heike Argstatter<sup>1</sup>, Sebastian Hoth<sup>2</sup>, Gerhard Dyckhoff<sup>2</sup>, Anne Kathrin Nickel<sup>1</sup>,  
Hagen Weidauer<sup>2</sup>, Hans Volker Bolay<sup>3</sup>

<sup>1</sup>Deutsches Zentrum für Musiktherapieforschung (German Center for Music Therapy Research)

<sup>2</sup>HNO Klinik, Universität Heidelberg (University Hospital for Ear, Nose and Throat, University Heidelberg)

<sup>3</sup>Fachhochschule Heidelberg (University of Applied Sciences Heidelberg)

## Background


- Tinnitus is one of the most common disorders in ENT medicine / otorhinolaryngology
- Patients suffering from chronic tinnitus experience psychologic distress such as sleeping disturbance, attention deficits, anxiety and depression
- Brain imaging (PET, fMRI, MEG) revealed plasticity in the auditory cortex similar to reorganization observed in chronic phantom pain
- Due to the complexity of the disease tinnitus, a comprehensive and interdisciplinary treatment is required

## Objectives

- Development and evaluation of a music therapeutic treatment concept for patients suffering from chronic, subjective, decompensated tinnitus
- Implementation of treatment standards

## Methods

- Randomized, prospective pilot study
- Sample: 20 patients (10 male, 10 female; mean age  $51 \pm 7$  yrs), suffering from decompensated chronic tinnitus were included in the study. All subjects underwent medical and psychological checkup in order to rule out organic and psychiatric diseases. Subsequently patients were randomly allocated to one of two groups: music therapy or control group



- Target variables: tinnitus variables (TQ, Goebel & Hiller, 1998), psychological factors
- Data collection: pre-post-measurements, follow-up after 24 weeks, for the music therapy group additional process measurement every 4 weeks


## Treatment

- All patients received comprehensive tinnitus counselling. The music therapy consisted of 12 sessions à 50 minutes individual music therapy. The control group did not obtain any further treatment
- Aims of the tinnitus therapy are the integration of the tinnitus into a music controllable hearing process and subsequently attenuation of subjective annoyance

## Results


### Tinnitus-Questionnaire over time

- Tinnitus pathology decreases continuously in the course of music therapy
- Regression analyses indicate a highly statistical and clinical significant decline in mean TQ-scores ( $B = -3,8$ ;  $F_{(1)} = 26,38$ ;  $p = .000$ )



### Group Comparison

- TQ-Scores diminish in the music therapy group by 24,9 points or 53% and in the control group by 2,4 points or 5%
- Results of ANOVA reveal significant group differences pre-post ( $F_{(4)} = 5,99$ ;  $p = .002$ ) with a large effect size of  $d = 1,73$



### Individual Changes

Clinical significant reduction ( <i>Winner</i> ) (-16)	N = 8	-30 points	-63%
Reliable reduction ( <i>Responder</i> ) (-6 to -15)	N = 1	-8 points	-20%
No change ( <i>Nonresponder</i> ) (-5 to +15)	N = 1	± 0 points	± 0%
Aggravation ( <i>Looser</i> ) (+16)	N = 0	---	---

## Discussion

Despite the small sample size, the innovative music therapy concept shows statistical and clinical significant results. Further research including brain imaging seems to be beneficial and is projected.